

On p -adic Twisted Euler (h, q) - l -Functions

S. H. Rim ^a, Y. Simsek ^b, V. Kurt ^b and T. Kim ^c

^aDepartment of Mathematics Education, Kyungpook National University,
Daegu 702-701, S. Korea

^bDepartment of Mathematics, Akdeniz University, 07058 Antalya, Turkey*

^cEECS, Kyungpook National University, Daegu 702-701, S. Korea
shrim@knu.ac.kr, ysimsek@akdeniz.edu.tr, vkurt@akdeniz.edu.tr, tkim@knu.ac.kr

Abstract : In the recent paper, Kim-Rim have studied interesting twisted q -Euler numbers and polynomials. In [8], Kim-Rim suggested the question to find a q -analogue of the p -adic twisted (h, q) - l -function which interpolates generalized twisted (h, q) -Euler numbers attached to χ . This question is remained open. The purpose of this paper is to give the answer of the question.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ are respectively denoted as the ring of p -adic rational integers, the field of p -adic rational numbers, the complex numbers field and the completion of algebraic closure of $\mathbb{Q}_p$. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{-v_p(p)} = \frac{1}{p}$.

When one talks of q -extension, q is considered in many ways such as an indeterminate, a complex number $q \in \mathbb{C}$, or p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$ one normally assumes that $|q| < 1$. If $q \in \mathbb{C}_p$, we normally assume that $|1 - q|_p < p^{-\frac{1}{p-1}}$, so that $q^x = \exp(x \log q)$ for $|x|_q \leq 1$.

We use the notations as

$$\begin{aligned} [x]_q &= \frac{1 - q^x}{1 - q} = 1 + q + q^2 + \cdots + q^{x-1}, \\ [x]_{-q} &= \frac{1 - (-q)^x}{1 + q} = 1 - q + q^2 - q^3 + \cdots + (-1)^x q^{x-1}. \end{aligned}$$

Let $UD(\mathbb{Z}_p)$ be the set of uniformly differentiable function on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, Kim originally defined the p -adic invariant q -integral on $\mathbb{Z}_p$ as follows:

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x, \text{ cf. [6], [7], [10],}$$

where N is natural number. Let

$$I_1(f) = \lim_{q \rightarrow 1} I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_1(x) = \lim_{N \rightarrow \infty} \frac{1}{p^N} \sum_{x=0}^{p^N-1} f(x),$$

*Second and third authors are supported by Akdeniz University Scientific Research Project Unit.

where N is a natural number (see [2], [3], [4], [5], [9]).

Let d be a fixed integer. For any positive integer N , we set

$$\begin{aligned}\mathbb{X} &= \mathbb{X}_d = \varprojlim_N (\mathbb{Z}/dp^N\mathbb{Z}), \\ \mathbb{X}^* &= \bigcup_{\substack{0 < a < dp \\ (a,p)=1}} (a + dp\mathbb{Z}_p), \\ a + dp^N &= \{x \in \mathbb{X} : x \equiv a \pmod{dp^N}\},\end{aligned}$$

where $a \in \mathbb{Z}$ lies in $0 \leq a < dp^N$.

Let us define $I_{-q}(f)$ as

$$I_{-q}(f) = \lim_{q \rightarrow -q} I_q(f) = \int_{\mathbb{Z}_p} f(x) d_{-q}(x) ..$$

This integral, $I_{-q}(f)$, can be considered as the q -deformed p -adic invariant integral on $\mathbb{Z}_p$ in the sense of fermionic, cf. [3], [4], [5], [6], [8].

In [6], multiple q -Euler polynomials of higher order were defined by

$$E_{n,q}^{(h,k)}(x) = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} [x + x_1 + \cdots + x_k]_q^n q^{\sum_{i=1}^k x_i(h-i)} d\mu_{-q}(x) \cdots d\mu_{-q}(x),$$

$k\text{-times}$

where $h \in \mathbb{Z}$, $k \in \mathbb{N}$. The q -Euler polynomials of higher order at $x = 0$ are called q -Euler numbers of higher order.

In [1], Carlitz originally constructed q -Bernoulli numbers and polynomials. These numbers and polynomials are studied by many authors (see [7], [8], [10], [11]). In particular, twisted (h, q) -Bernoulli numbers and polynomials were also studied by several authors (see [9], [10], [11]).

In [8], Kim-Rim introduced an interesting twisted q -Euler numbers and polynomials associated with basic twisted q - l -functions and suggested the following question:

“Find a q -analogue of the p -adic twisted l -function which interpolates generalized twisted q -Euler numbers attached to χ , $E_{n,w,\chi,q}$.”

In this paper, we give some interesting properties related to twisted (h, q) -Euler numbers and polynomials. The final purpose of this paper is to construct p -adic twisted Euler (h, q) - l -function which is a part of answer for the question in [8].

2. p -adic Invariant Integral on $\mathbb{Z}_p$ Associated with Twisted (h, q) -Euler Numbers and Polynomials

Let $h \in \mathbb{Z}$ and $q \in \mathbb{C}_p$ with $|1 - q|_p < p^{-\frac{1}{p-1}}$. From the invariant integral on $\mathbb{Z}_p$ in the sense of fermionic, we define

$$I_{-1}(f) = \lim_{q \rightarrow -1} I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x),$$

where $f \in UD(\mathbb{Z}_p)$, cf. [3]. Note that $I_{-1}(f_1) + I_{-1}(f) = 2f(0)$, where $f_1(x) = f(x+1)$. Let $C_{p^n} = \{\xi : \xi^{p^n} = 1\}$ be the cyclic group of order p^n and let $T_p = \lim_{n \rightarrow \infty} C_{p^n} = C_{p^\infty}$. Then T_p is

p -adic locally constant space. For $\xi \in T_p$, we denote by $\phi_\xi : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ defined by $\phi_\xi(x) = \xi^x$ be the locally constant function. If we take $f(x) = \phi_\xi(x) e^{tx}$, then we have

$$\int_{\mathbb{Z}_p} e^{tx} \phi_\xi(x) d\mu_{-1}(x) = \frac{2}{\xi e^t + 1}, \text{ (see [3])}. \quad (1)$$

In complex case the twisted Euler numbers were defined by Kim-Rim [8]

$$\frac{2}{\xi e^t + 1} = \sum_{n=0}^{\infty} E_{n,\xi} \frac{t^n}{n!},$$

where $|\log \xi + t| < \pi$. Thus we have

$$\int_{\mathbb{Z}_p} x^n \phi_\xi(x) d\mu_{-1}(x) = E_{n,\xi}, \quad (n \geq 0).$$

By using iterative method of p -adic invariant integral on $\mathbb{Z}_p$ in the sense of fermionic, we see that

$$I_{-1}(f_n) = (-1)^n I_{-1}(f) + 2 \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l), \quad (2)$$

where $f_n(x) = f(x+n)$. If n is odd positive integer, then we have

$$I_{-1}(f_n) + I_{-1}(f) = 2 \sum_{l=0}^{n-1} (-1)^l f(l). \quad (3)$$

Let χ be the Dirichlet's character with conductor d ($=\text{odd}$) $\in \mathbb{N}$, and let $f(x) = \chi(x) \phi_\xi(x) e^{tx} \in UD(\mathbb{Z}_p)$. From (3), we can derive the following:

$$\int_{\mathbb{X}} e^{tx} \phi_\xi(x) \chi(x) d\mu_{-1}(x) = 2 \frac{\sum_{a=0}^{d-1} \chi(a) \phi_\xi(x) e^{at}}{\xi^d e^{dt} + 1}. \quad (4)$$

Now we define twisted generalized Euler numbers attached to χ as follows:

$$\frac{2 \sum_{a=0}^{d-1} \chi(a) \phi_\xi(x) e^{at}}{\xi^d e^{dt} + 1} = \sum_{n=0}^{\infty} E_{n,\chi,\xi} \frac{t^n}{n!}.$$

From (1) and (4), we note that

$$\begin{aligned} \int_{\mathbb{Z}_p} x^n \phi_\xi(x) d\mu_{-1}(x) &= E_{n,\xi}, \\ \int_{\mathbb{X}} x^n \phi_\xi(x) \chi(x) d\mu_{-1}(x) &= E_{n,\chi,\xi} \end{aligned} \quad (5)$$

In [6], (h, q) -Euler numbers were defined by

$$E_{n,q}^{(h,1)}(x) = \int_{\mathbb{Z}_p} q^{(h-1)y} [x+y]_q^n d\mu_{-q}(y),$$

where $h \in \mathbb{Z}$. $E_{n,q}^{(h,1)}(0) = E_{n,q}^{(h,1)}$ will be called (h, q) -Euler number. In the special case $h = 1$, we note that $\lim_{h \rightarrow 1} E_{n,q}^{(h,1)} = E_{n,q}^{(1,1)}$ becomes q -Euler numbers, which were originally defined by Kim [6]. That is $\lim_{h \rightarrow 1} E_{n,q}^{(h,1)} = E_{n,q}^{(1,1)} = E_{n,q}$.

In the viewpoint of (5), we consider twisted (h, q) -Euler numbers using p -adic invariant q -integral on $\mathbb{Z}_p$ in the sense of fermionic as follows:

$$E_{n,\xi,q}^{(h,1)}(x) = \int_{\mathbb{Z}_p} q^{(h-1)y} \phi_\xi(y) [x+y]_q^n d\mu_{-q}(y), \quad (6)$$

which are called twisted (h, q) -Euler polynomials. In the special case $x = 0$, we use notation $E_{n,\xi,q}^{(h,1)}(0) = E_{n,\xi,q}^{(h,1)}$ which are called (h, q) -twisted Euler numbers. Note that

$$E_{n,\xi,q}^{(h,1)}(x) = \frac{[2]_q}{(1-q)^n} \sum_{j=0}^n \binom{n}{j} (-1)^j q^{xj} \frac{1}{1 + \xi q^{h+j}}, \quad (7)$$

where

$$\binom{n}{j} = \frac{n(n-1) \cdots (n-j+1)}{j!}.$$

From (7), we note that

$$E_{n,\xi,q}^{(h,1)}(x) = [2]_q \sum_{k=0}^{\infty} (-1)^k \xi^k q^{hk} [x+k]_q^n, \quad (8)$$

where $h \in \mathbb{Z}$, $n \in \mathbb{N}$. Equation (8) is equivalent to

$$E_{n,\xi,q}^{(h,1)}(x) = \frac{[2]_q}{[2]_{q^d}^n} [d]_q^n \sum_{a=0}^{d-1} (-1)^a \xi^a q^{ha} E_{n,\xi^d,q^d}^{(h,1)}\left(\frac{x+a}{d}\right)$$

(distribution for $E_{n,\xi,q}^{(h,1)}(x)$), where $n, d (= \text{odd}) \in \mathbb{N}$.

Let χ be the Dirichlet character with conductor $f (= \text{odd}) \in \mathbb{N}$. Then we define the generalized twisted (h, q) -Euler numbers attached to χ as follows: For $n \geq 0$,

$$E_{n,\xi,\chi,q}^{(h,1)} = \int_{\mathbb{X}} \chi(x) q^{(h-1)x} \xi^x [x]_q^n d\mu_{-q}(x), \quad (9)$$

where $h \in \mathbb{Z}$. Note that $E_{n,1,\chi,q}^{(1,1)} = E_{n,\chi,q}$, see [3]. From (9), we also derive

$$\begin{aligned} E_{n,\xi,\chi,q}^{(h,1)} &= [f]_q^n \frac{[2]_q}{[2]_{q^f}^n} \sum_{a=0}^{f-1} \chi(a) (-1)^a \xi^a q^{ha} E_{n,\xi^f,q^f}^{(h,1)}\left(\frac{a}{f}\right) \\ &= [2]_q \sum_{k=1}^{\infty} \chi(k) (-1)^k \xi^k q^{hk} [k]_q^n, \end{aligned} \quad (10)$$

where $n, d (= \text{odd}) \in \mathbb{N}$.

3. Twisted (h, q) -Euler Zeta Function in $\mathbb{C}$

For $q \in \mathbb{C}$ with $|q| < 1$, $s \in \mathbb{C}$, we define

$$\zeta_{E,q,\xi}^{(h,1)}(s) = [2]_q \sum_{k=1}^{\infty} \frac{(-1)^k \xi^k q^{hk}}{[k]_q^s}.$$

Then we see that $\zeta_{E,q,\xi}^{(h,1)}(s)$ is analytic continuation in whole complex plane. We easily see that

$$\zeta_{E,q,1}^{(h,1)}(s) = \zeta_{E,q}^{(h,1)}(s), \text{ see [6].}$$

We now also consider Hurwitz's type twisted (h, q) -Euler zeta function as follows: For $s \in \mathbb{C}$, define

$$\zeta_{E,q,\xi}^{(h,1)}(s, x) = [2]_q \sum_{k=0}^{\infty} \frac{(-1)^a \xi^k q^{hk}}{[x+k]_q^s}, \quad (11)$$

for $s \in \mathbb{C}$, $h \in \mathbb{Z}$. By (8) and (11), we see that

$$\zeta_{E,q,\xi}^{(h,1)}(-n, x) = E_{n,\xi,q}^{(h,1)}(x),$$

where $n \in \mathbb{N}$, $h \in \mathbb{Z}$.

Let χ be the Dirichlet's character with conductor $f (= \text{odd}) \in \mathbb{N}$. Then we define twisted (h, q) - l -function which interpolates twisted generalized (h, q) -Euler numbers attached to χ as follows: For $s \in \mathbb{C}$, $h \in \mathbb{Z}$, we define

$$l_{q,\xi}^{(h,1)}(s, \chi) = [2]_q \sum_{k=1}^{\infty} \frac{\chi(k) (-1)^k q^{hk} \xi^k}{[k]_q^s}. \quad (12)$$

For any positive integer n , we have

$$l_{q,\xi}^{(h,1)}(-n, \chi) = E_{n,\xi,\chi,q}^{(h,1)}, \quad n \in \mathbb{N}. \quad (13)$$

From (13), we derive

$$\begin{aligned} l_{q,\xi}^{(h,1)}(s, \chi) &= [2]_q \sum_{k=1}^{\infty} \frac{\chi(k) (-1)^k q^{hk} \xi^k}{[k]_q^s} \\ &= [f]_q^{-s} \frac{[2]_q}{[2]_{q^f}} \sum_{a=1}^f \chi(a) (-1)^a \xi^a q^{ha} \zeta_{E,\xi^f,q^f}^{(h,1)}\left(s, \frac{a}{f}\right). \end{aligned}$$

Let s be a complex variable and let a and $F (= \text{odd})$ be integer with $0 < a < F$. We consider the following twisted (h, q) -harmonic sums (or partial (h, q) -zeta function):

$$\begin{aligned} H_{E,q,\xi}^{(h,1)}(s, a|F) &= \sum_{\substack{m \equiv 0 \pmod{F} \\ m > 0}} \frac{(-1)^m q^{hm} \xi^m}{[m]_q^s} \\ &= \sum_{n=0}^{\infty} \frac{(-1)^{a+nF} q^{h(a+nF)} \xi^{a+nF}}{[a+nF]_q^s} \\ &= (-1)^a q^{ha} \xi^a \sum_{n=0}^{\infty} \frac{(-1)^n (q^F)^{hn} (\xi^F)^n}{[F]_q^s \left[\frac{a}{F} + n\right]_{q^F}^s} \\ &= [F]_q^{-s} \frac{(-1)^a q^{ha} \xi^a}{[2]_{q^F}} \zeta_{E,\xi^F,q^F}^{(h,1)}\left(s, \frac{a}{F}\right). \end{aligned}$$

Thus we have

$$H_{E,q,\xi}^{(h,1)}(s, a|F) = [F]_q^{-s} \frac{(-1)^a q^{ha} \xi^a}{[2]_{q^F}} \zeta_{E,\xi^F,q^F}^{(h,1)}\left(s, \frac{a}{F}\right). \quad (14)$$

By (12), (13) and (14), we see that

$$l_{q,\xi}^{(h,1)}(s, \chi) = [2]_q \sum_{a=1}^F \chi(a) H_{E,q,\xi}^{(h,1)}(s, a|F). \quad (15)$$

From (14), we note that

$$H_{E,q,\xi}^{(h,1)}(-n, a|F) = \frac{[F]_q^n}{[2]_{q^F}} (-1)^a q^{ha} \xi^a E_{n,\xi^F,q^F}^{(h,1)}\left(\frac{a}{F}\right), \quad (16)$$

where n is a positive integer. By (10), (15) and (16), we see that

$$l_{q,\xi}^{(h,1)}(-n, \chi) = E_{n,\xi,\chi,q}^{(h,1)}.$$

The Euler (h, q) -twisted harmonic sum $H_{E,q,\xi}^{(h,1)}(s, a|F)$ will be called partial twisted (h, q) -zeta function which interpolates twisted (h, q) -Euler polynomials at negative integers. The values $l_{q,\xi}^{(h,1)}(s, \chi)$ are algebraic, hence regarded as lying in the extension of $\mathbb{Q}_p$.

4. p -adic Twisted Euler (h, q) - l -Function

Let $\omega(x)$ be the Teichmüller character and let $\langle x \rangle_q = \langle x \rangle = \frac{[x]_q}{\omega(x)}$. When $F (= \text{odd})$ is multiple of p , and $(a, p) = 1$, we define p -adic partial (h, q) -zeta function as follows: For $h \in \mathbb{Z}$, $q \in \mathbb{C}_p$ with $|1 - q|_p < p^{-\frac{1}{p-1}}$, we define

$$H_{E,p,q,\xi}^{(h,1)}(s, a|F) = \frac{(-1)^a \xi^a q^{ha}}{[2]_{q^F}} \langle a \rangle^{-s} \sum_{j=0}^{\infty} \binom{-s}{j} \left(\frac{[F]_q}{[a]_q} \right)^j q^{aj} E_{j,\xi^F,q^F}^{(h,1)},$$

where $s \in \mathbb{Z}_p$. Thus we note that

$$\begin{aligned} H_{E,p,q,\xi}^{(h,1)}(-n, a|F) &= \frac{(-1)^a \xi^a q^{ha}}{[2]_{q^F}} \langle a \rangle^n \sum_{j=0}^n \binom{n}{j} \left(\frac{[F]_q}{[a]_q} \right)^j q^{aj} E_{j,\xi^F,q^F}^{(h,1)} \\ &= \frac{(-1)^a \xi^a q^{ha}}{[2]_{q^F}} [F]_q^n \omega^{-n}(a) \sum_{j=0}^n \binom{n}{j} \left(\frac{[a]_q}{[F]_q} \right)^{n-j} (q^F)^{\frac{a}{F}j} E_{j,\xi^F,q^F}^{(h,1)} \\ &= [F]_q^n \frac{(-1)^a \xi^a q^{ha}}{[2]_{q^F}} \omega^{-n}(a) E_{n,\xi^F,q^F}^{(h,1)}\left(\frac{a}{F}\right) \\ &= \omega^{-n}(a) H_{E,q,\xi}^{(h,1)}(-n, a|F). \end{aligned}$$

Therefore, we obtain the following formula:

$$H_{E,p,q,\xi}^{(h,1)}(-n, a|F) = \omega^{-n}(a) H_{E,q,\xi}^{(h,1)}(-n, a|F). \quad (17)$$

Now we consider p -adic interpolating function for twisted generalized (h, q) -Euler numbers attached to χ as follows:

$$l_{p,q,\xi}^{(h,1)}(s, \chi) = [2]_q \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) H_{E,p,q,\xi}^{(h,1)}(s, a|F), \quad (18)$$

for $s \in \mathbb{Z}_p$. Let n be a natural number. Then we have

$$\begin{aligned}
l_{p,q,\xi}^{(h,1)}(-n, \chi) &= [2]_q \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) H_{E,p,q,\xi}^{(h,1)}(-n, a|F) \\
&= [2]_q \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) [F]_q^n \frac{(-1)^a \xi^a q^{ha}}{[2]_{q^F}} \omega^{-n}(a) E_{n,\xi^F,q^F}^{(h,1)}\left(\frac{a}{F}\right) \\
&= [F]_q^n \frac{[2]_q}{[2]_{q^F}} \sum_{\substack{a=1 \\ (a,p)=1}}^F (-1)^a \chi \omega^{-n}(a) \xi^a q^{ha} E_{n,\xi^F,q^F}^{(h,1)}\left(\frac{a}{F}\right) \\
&= E_{n,\xi,\chi \omega^{-n},q}^{(h,1)} - \chi \omega^{-n}(p) [p]_q^n \frac{[2]_q}{[2]_{q^p}} E_{n,\xi^p,\chi \omega^{-n},q^p}^{(h,1)}.
\end{aligned}$$

Therefore we obtain the following theorem:

Theorem : For $s \in \mathbb{Z}_p$, we define p -adic twisted Euler (h, q) - l -function as follows:

$$\begin{aligned}
l_{p,q,\xi}^{(h,1)}(s, \chi) &= [2]_q \sum_{\substack{a=1 \\ (a,p)=1}}^F \chi(a) H_{E,p,q,\xi}^{(h,1)}(s, a|F) \\
&= [2]_q \sum_{\substack{k=1 \\ (k,p)=1}}^{\infty} \frac{(-1)^k \chi(k) q^{hk} \xi^k}{[k]_q^s}.
\end{aligned}$$

Then we have

$$l_{p,q,\xi}^{(h,1)}(-n, \chi) = E_{n,\xi,\chi \omega^{-n},q}^{(h,1)} - \chi \omega^{-n}(p) [p]_q^n \frac{[2]_q}{[2]_{q^p}} E_{n,\xi^p,\chi \omega^{-n},q^p}^{(h,1)}.$$

Remark : From the above theorem, we note that

$$l_{p,q,\xi}^{(h,1)}(s, \chi) = \int_{\mathbb{X}^*} \chi(x) \langle x \rangle^{-s} q^{(h-1)x} \xi^x d\mu_{-q}(x).$$

References

- [1] L. Carlitz, q -Bernoulli and Eulerian numbers, Trans. Amer. Math. Soc. 76 (1954), 332-370.
- [2] M. Cenkci, M. Can, Some results on q -analogue of the Lerch zeta function, Advan. Stud. Contemp. Math. 12 (2006), 213-223.
- [3] T. Kim, On the analogs of Euler numbers and polynomials associated with p -adic q -integral on $\mathbb{Z}_p$ at $q = -1$, J. Math. Anal. Appl. (2006), doi:10.1016/j.jmaa.2006.09.029.
- [4] T. Kim, On p -adic q - l -functions and sums of powers, J. Math. Anal. Appl. (2006), doi:10.1016/j.jmaa.2006.07.071.
- [5] T. Kim, On the q -extension of Euler and Genocchi numbers, J. Math. Anal. Appl. 326 (2007), 1458-1469.
- [6] T. Kim, q -Volkenborn integration and applications, arXiv.math.NT/0510524.

- [7] T. Kim, S. H. Rim, q -Volkenborn integration, Russian J. Math. Phys. 9 (2002), 288-299.
- [8] T. Kim, S. H. Rim, On the twisted q -Euler numbers and polynomials associated with basic q - l -functions, arXiv.math.NT/0611807.
- [9] T. Kim, L. C. Jang, S. H. Rim, H. K. Pak, On the twisted q -zeta functions and q -Bernoulli polynomials, Far East J. Appl. Math. 13 (2003), 13-21.
- [10] Y. Simsek, Twisted (h, q) -Bernoulli numbers and polynomials related to twisted (h, q) -zeta function and L -function, J. Math. Anal. Appl. 324 (2006), 790-804.
- [11] Y. Simsek, On p -adic twisted q - L -functions related to generalized twisted Bernoulli numbers, Russian J. Math. Phys. 13 (3) (2006), 340-348.